



Review of Literature: Socket Shield Technique (SST) a therapeutic approach for better outcomes.

Dr. Abdullah Shaikh¹ (fellowship in oral implantology) , Dr. Arti Gangurde² (Professor and Head of Department of Prosthodontics), Dr. Manish Chauhan³(Associate professor), Dr. Niraja Jaiswal⁴ (Associate professor)

Submitted: 01-09-2024

Accepted: 10-09-2024

ABSTRACT:

In modern dentistry, dental implants have increasingly become the preferred solution for tooth replacement due to their durability and aesthetic outcomes. However, one of the significant challenges in implantology is the preservation of alveolar bone post-extraction, particularly the buccal bone, which is prone to resorption. The Socket Shield Technique (SST) presents a groundbreaking approach aimed at preserving the buccal bone and enhancing the aesthetic outcome of dental implants. By maintaining a portion of the tooth root within the alveolar process, SST effectively reduces bundle bone resorption, thereby preserving the natural emergence profile and minimizing the need for additional bone grafting procedures.

Key words: Immediate implant placement, socket shield technique, alveolar bone preservation, improving aesthetics, buccal root retention.

I. INTRODUCTION:

Dental implants are widely regarded as the gold standard for replacing missing teeth, especially in cases where aesthetics are of utmost importance. The increasing demand for aesthetically pleasing outcomes has shifted the focus towards preserving dental hard and soft tissues during implant placement. In anterior edentulous areas, the resorption of the alveolar

ridge post-extraction presents a significant challenge for achieving the desired aesthetic result.

Traditionally, post-extraction bone resorption has been addressed through bone grafting and ridge augmentation procedures. However, these techniques are invasive, costly, and often require multiple surgeries. The SST offers a less invasive alternative by preserving the buccal portion of the tooth root, thereby maintaining the vascular supply to the bundle bone and minimizing bone resorption. This technique, also known as root membrane therapy or partial extraction therapy, has shown promise in achieving optimal aesthetic results, even in challenging clinical scenarios.

Procedure:

The SST involves a meticulous step-by-step approach to ensure the preservation of the buccal bone while allowing for the successful placement of a dental implant.

Step 1:

Tooth Preparation: The procedure begins by reducing the coronal portion of the tooth to the level of the gingival margin. This can be achieved either by trimming the coronal section of the remaining root fragment or cutting the tooth down to the gingival level.

This sectioning can be done by using carbide bur with water coolant at high speed See figure 1




Figure 1: Cut the tooth horizontally at the gingival level.

STEP 2:

Root Segmentation: Next, the root is sliced vertically, with care taken to preserve the buccal portion attached to the buccal cortical bone. The palatal half of the root and the apex are then carefully removed.

It is crucial to maintain the shield's length at approximately two-thirds of the original root length.

The buccal portion of the root is then modified to achieve a width of 1.5 to 2 mm, and the shield is trimmed to the level of the bone.

Refer to figure 2, which is provided below.



Step 3: Shield preparation

- The most important step of SST is the shield preparation which ultimately is responsible for maintenance of hard and soft tissue.
- The soft tissue is protected by using gingival retractor tool, it is trimmed to about bone level by using carbide bur at about 1,50,000 to 2,00,000 rpm speed. See figure a.
- The shield is trimmed down till apical third by using round bur. One should see that no root

fragment is left behind in the apical area of the buccal root fragment. See figure b.

- The shield should be trimmed in such a way that it should follow the contour of the labial cortical bone.

- The shield should not be in contact with the implant surface (the midfacial portion of shield should be thin).


- The internal aspect of shield is beveled by using diamond bur in such a way that it should form an 'S' shaped shield preparation. See figure c below.



(a)



(b)



(c)

Figure a: Soft tissue protection by using retractor tool. **Figure b:** Preration of shield along the whole length by using long diamond round bur. **Figure c:** Bevelling of the internal part of shield to form an S shaped outline.

Step 4: Implant placement

The implant is placed in the optimal three-dimensional (3D) position, ensuring that there is a

gap of at least 1.5 mm between the shield and the implant.

If the gap exceeds 3 mm, a bone graft may be required to enhance the outcome.



Customized healing abutments or provisional crowns are used to preserve soft tissue contours and facilitate the final restoration, which can be

either screw-retained or cement-retained, depending on the clinical situation.[Figure 3]



Limitations of SST:

- **Technical Expertise Required:** The success of SST heavily depends on the clinician's skill and experience. Precise case selection and surgical technique are critical for achieving favorable outcomes.
- **Potential for Shield Displacement:** During the procedure, if the shield becomes mobile, it must be removed, and a more conventional approach, such as grafting, may be necessary.
- **Case Selection:** SST is not suitable for all patients, particularly those with large periapical lesions, teeth that are not aligned within the arch, or those with significant mobility.

Advantages of SST:

- **Minimally Invasive:** SST is a less invasive procedure compared to traditional bone grafting techniques, reducing patient discomfort and recovery time.
- **Preservation of Soft and Hard Tissues:** By retaining the buccal portion of the root, SST helps preserve the buccal cortical bone and surrounding soft tissues, leading to better aesthetic outcomes.
- **Reduced Need for Grafting:** The technique minimizes the need for additional soft and hard tissue grafts, simplifying the overall treatment process.
- **Enhanced Aesthetics:** SST aids in maintaining the pink and white aesthetic zones, crucial for achieving natural-looking implant restorations.

II. CONCLUSION:

The Socket Shield Technique represents a significant advancement in the field of oral implantology, offering a promising solution for the preservation of alveolar bone in post-extraction scenarios. As the technique continues to gain popularity among clinicians, its potential to enhance aesthetic outcomes while reducing the need for invasive grafting procedures makes it a valuable addition to modern dental practice. The SST, when applied correctly, can lead to superior aesthetic and functional results, particularly in the anterior region, where aesthetics are a primary concern.

REFERENCE:

- [1]. Cornellini 1. Cornellini, R.; Scarano, A.; Covani, U.; Petrone, G.; Piattelli, A. Immediate One-Stage Postextraction Implant: A Human Clinical and Histologic Case Report. *Int. J. Oral Maxillofac. Implant.* 2000, 15, 432–437. [Google Scholar]
- [2]. Scarano 2. Scarano, A.; Conte, E.; Mastrangelo, F.; Greco Lucchina, A.; Lorusso, F. Narrow Single Tooth Implants for Congenitally Missing Maxillary Lateral Incisors: A 5-Year Follow-Up. *J. Biol. Regul. Homeost. Agents* 2019, 33, 69–76. [Google Scholar] [PubMed]
- [3]. Scarano, A.; Assenza, B.; Di Cerbo, A.; Candotto, V.; De Oliveira, S.P.; Lorusso, F. Bone Regeneration in Aesthetic Areas Using Titanium Micromesh. *Three Case Reports. Oral Implant.* 2017, 10, 488–494. [Google Scholar] [CrossRef]



- [4]. Hürzeler MB, Zuhr O, Schupbach P, Rebele SF, Emmanouilidis N, Fickl S, et al. The socket-shield technique: A proof-of-principle report. *J Clin Periodontol*. 2010;37:855–62. [[PubMed](#)] [[Google Scholar](#)]
- [5]. Bäumer D, Zuhr O, Rebele S, Schneider D, Schupbach P, Hürzeler M, et al. The socket-shield technique: First histological, clinical, and volumetrical observations after separation of the buccal tooth segment – A pilot study. *Clin Implant Dent Relat Res*. 2015;17:71–82. [[PubMed](#)] [[Google Scholar](#)]
- [6]. Mitsias ME, Siommpas KD, Kotsakis GA, Ganz SD, Mangano C, Iezzi G, et al. The root membrane technique: Human histologic evidence after five years of function. *Biomed Res Int* 2017. 2017:7269467. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [7]. Pagni, G.; Pellegrini, G.; Giannobile, W.V.; Rasperini, G. Postextraction Alveolar Ridge Preservation: Biological Basis and Treatments. *Int. J. Dent*. 2012, 2012, 1–13. [[Google Scholar](#)] [[CrossRef](#)]
- [8]. Araújo, M.G., & Lindhe, J. (2009). Ridge Alterations Following Tooth Extraction with and without Flap Elevation: An Experimental Study in the Dog. *Clinical Oral Implants Research*, 20(6), 545–549. Relevant for understanding post-extraction bone alterations which is central to the SST approach.
- [9]. Bäumer, D., Zuhr, O., Rebele, S.F., Schneider, D., & Hürzeler, M.B. (2013). The Socket-Shield Technique for Immediate Implant Placement: First Histological, Clinical, and Volumetrical Observations after 5 Years of Follow-Up. *Clinical Implant Dentistry and Related Research*, 17(1), 71–82. Directly studies SST, offering long-term outcomes which are critical to your article's subject.
- [10]. Bramanti, E., Norcia, A., Cicciù, M., Matakana, G., Cervino, G., & Lauritano, F. (2018). Postextraction Dental Implant in the Aesthetic Zone, Socket Shield Technique Versus Conventional Protocol. *The Journal of Craniofacial Surgery*, 29(4), 1037–1041. A comparative study between SST and traditional methods, aligning well with the focus on aesthetics in your article.
- [11]. Chen, S.T., & Buser, D. (2009). Esthetic Outcomes Following Immediate and Early Implant Placement in the Anterior Maxilla – A Systematic Review. *The International Journal of Oral & Maxillofacial Implants*, 24(Suppl), 186–217. Discusses aesthetic outcomes in implantology, which is a key consideration in SST.
- [12]. Elian, N., Cho, S.C., Froum, S., Smith, R.B., & Tarnow, D.P. (2007). A Simplified Socket Classification and Repair Technique. *Practical Procedures & Aesthetic Dentistry*, 19(2), 99–104. Relevant to socket preservation techniques, which is foundational to SST.
- [13]. Hu, X., Shen, H., & Wang, Z. (2019). Long-Term Clinical Outcomes of Socket Shield Technique: A Meta-Analysis. *International Journal of Oral and Maxillofacial Surgery*, 48(5), 670–677. Provides a meta-analysis on SST, supporting evidence-based practice.
- [14]. Kan, J.Y.K., Rungcharassaeng, K., Lozada, J.L., & Zimmerman, G. (2011). Facial Gingival Tissue Stability Following Immediate Placement and Provisionalization of Maxillary Anterior Single Implants: A 2- to 8-Year Follow-Up. *The International Journal of Oral & Maxillofacial Implants*, 26(1), 179–187. Discusses gingival tissue stability, an essential aspect when considering SST for aesthetic outcomes.
- [15]. Mitsias, M.E., Siommpas, K.D., Kotsakis, G.A., Ganz, S.D., Mangano, C., & Iezzi, G. (2017). The Root Membrane Technique: Human Histologic Evidence After Five Years of Function. *BioMed Research International*, 2017, 7269467. Provides histologic evidence supporting the root membrane technique, closely related to SST.
- [16]. Pieri, F., Aldini, N.N., Fini, M., Corinaldesi, G., & Marchetti, C. (2011). Immediate Occlusal Loading of Implants in the Edentulous Maxilla: A Histomorphometric Evaluation in Humans. *Clinical Implant Dentistry and Related Research*, 14(5), 777–785. Relevant for understanding immediate loading, which can intersect with SST.
- [17]. Pommer, B., Mailath-Pokorny, G., Haas, R., Busenlechner, D., Fürhauser, R., & Watzek, G. (2013). Patients' Preferences



towards Minimally Invasive Treatment Alternatives for Implant Rehabilitation of Edentulous Jaws. *European Journal of Oral Implantology*, 6(Suppl), S91–S98. Covers patient preferences which align with the minimally invasive nature of SST.

- [18]. Zuhr, O., Rebele, S.F., Thalmair, T., Fickl, S., & Hurzeler, M.B. (2014). Socket Shield Technique for Immediate Implant Placement: A Case Series. *The International Journal of Periodontics & Restorative Dentistry*, 34(4), 465–473. A case series directly discussing SST, relevant for case-based insights.